

Persantin Myocardial Perfusion Imaging for The Detection of Ischemia in Asymptomatic Diabetic Saudi Patients (DIADS)

Saleh Othman*

Department of Radiology and Medical Imaging, King Khalid University Hospital and College of Medicine, King Saud University, Riyadh, Saudi Arabia

*Corresponding author: Saleh Othman, Associate Professor and Consultant, Nuclear Medicine Division, Department of Radiology and Medical Imaging, King Khalid University Hospital and College of Medicine, King Saud University, Riyadh, Saudi Arabia, Tel: 00 966 1 4671159, E-mail: sothman@ksu.edu.sa

Received Date: January 15, 2024 Accepted Date: February 15, 2024 Published Date: February 18, 2024

Citation: Saleh Othman (2024) Persantin Myocardial Perfusion Imaging for The Detection of Ischemia in Asymptomatic Diabetic Saudi Patients (DIADS). J Radiol Nucl Med 1: 1-10.

Abstract

Background: Vascular complications occur in diabetic patients and more frequent when diabetes mellitus (DM) is uncontrolled. Among these complications are coronary artery diseases, which may lead to asymptomatic ischemic heart disease (IHD). The aim of this pilot study is to determine the prevalence of silent ischemia among Saudi patients with type 2 DM.

Results: A retrospective cohort study analyzing myocardial perfusion imaging (MPI) in Saudi patients with type 2 DM investigated for the presence of silent IHD. They had DM at least for 5 years duration and not less than 2 cardiovascular risk factors with no signs, symptoms or history of IHD. The mean age of the total cohort (n=490) was 62.2±11 years, 299 females (61%) and 191 males (39). Persantin MPI was positive in 237 patients (48.4%) and only 158 of them had coronary angiography of whom 33 (21%) had significant coronary anatomic lesions.

Conclusion: The prevalence of silent ischemia among asymptomatic Saudi patients with type 2 DM was 21%, which means that more than 2 out of 10 patients had silent myocardial ischemia. However and in order to justify the use of stress MPI as a screening test in this group of patients, a prospective large national study including both controlled and non-controlled type 2 DM is required.

Keywords: Saudi; Diabetics; Asymptomatic; Silent Ischemia; DIADS

List of Abbreviations: MIBI: technetium (99mTc) sestamibi; SSS: summed stress score; SRS: summed rest score; SDS: summed difference score; TID: transient ischemic dilation; LVEF: left ventricular ejection fraction

Introduction

Diabetes mellitus (DM) is a major cause of cardiovascular morbidity and mortality worldwide [1]. The prevalence of DM worldwide has been estimated to be 200 million as of 2007 and to 300 million in 2025 [2]. In a national epidemiological survey conducted in Saudi Arabia between 1995 and 2000, the prevalence of DM was 23.7%, one of the highest in the world [3]. Diabetic patients have more extensive atherosclerosis and a higher prevalence of multi-vessel coronary artery disease (CAD), as well as higher silent myocardial ischemia and infarction rates, than non-diabetics [4]. CAD is the leading cause of death among type 2 DM patients [1,5-8]. More than 50% of diabetics die of coronary artery disease [9]. The risk of cardiac death in patients with type 2 DM reported to be 2-4 times than non-diabetic patients [10-12]. Silent ischemia in these patient is of asymptomatic nature and clinical manifestation only discovered in advanced stage [10-13]. Silent ischemia reported to be as high as 22% in diabetics, with marked perfusion abnormalities in 6% of patients [14,15].

Myocardial perfusion imaging (MPI) is an accurate diagnostic tool for detecting the functional significance of obstructive CAD and estimating the burden of myocardial ischemia [16]. The sensitivity and specificity of MPI in detecting CAD reported to be 86% and 74% respectively [11,15,16].

The present study aimed to determine the prevalence of silent ischemia in asymptomatic Saudi patients with type 2 DM using pharmacologic stress MPI.

Methods

Study population

This retrospective study included consecutive patients previously diagnosed with type 2 DM referred to King Khalid University Hospital (Riyadh –Saudi Arabia) from the diabetic center at King Abdel Aziz University Hospital (Riyadh –Saudi Arabia) for stress MPI to investigate a suspicion of silent myocardial ischemia. We reviewed studies of all patients performed between July 2010 and October 2015. In addition, each patient had completed the full pharmacologic stress protocol and had interpretable imaging data. We used the patients' medical records to obtain demographic data, medical history, medical therapies, and subsequent cardiac procedures.

The study population consisted of 490 patients; 299 females (61%) and 191 males (39%) asymptomatic Saudi patients

with type 2 DM who had controlled DM for more than 5 years and at least two other risk factors such as hypertension and dyslipidemia. All patients had no symptoms or signs of ischemic heart disease. Patient age ranged from 40 to 70 years, 70 patients were less than 50 years old (14.3%) and 420 patients were more than 50 years old (85.7%).

Myocardial perfusion imaging protocol

All patients were instructed to stop their anti-angina medications, including β -blockers, calcium channel blockers, and long-acting nitrates, 48 hours prior to the stress date. The patients also asked to abstain from caffeinated beverages for 24 hours and to fast 6 hours prior to the test.

All patients underwent a one-day imaging protocol. Single photon emission computed tomography (SPECT) scan acquired as per the guidelines of the American Society of Nuclear Cardiology [17-19]. Patients underwent an intravenous weight-based dipyridamole (Persantine) infusion administered over 4 minutes. Vital signs (Blood pressure, heart rate) and ECG monitoring supervised by cardiology fellow during stress part of the study. The radioactive isotope used was technetium 99m-sestamibi. The radioactive isotope was injected at minute 4 of Persantin infusion. Technetium 99m-sestamibi (10 mCi) administered at the stress part of the study (The dose increased in overweight patients) done first, followed by 30 mCi given at rest. SPECT imaging was performed using a dual head gamma camera (Cardio MD, Phillips) for slim patients with additional prone images when diaphragmatic or breast artifact was there and images analyzed with AutoSPECT professional software without motion correction or attenuation correction software and SPECT low dose CT (Philips' BrightView XCT) for obese patients with attenuation correction. Butterworth filter and reconstruction using the filtered back projection method in all studies. Astonish reconstruction was used when time of imaging planned to be reduced due to patient condition or technical issues. Gated acquisition performed on the stress images in all patients except those with significant arrhythmia and high rejection rate.


Statistical analysis: Categorical data summarized as absolute numbers and percentages \pm standard deviation (SD).


Results

Table 1 depicts the baseline characteristics of the study population.

Table 1: Patient characteristics of the study group

Clinical Characteristic	No.	%
Gender: Male	191	39%
Female	299	61%
Age: <50	70	14.3%
>50	420	85.7%
Weight: Normal (BMI <25.0)	38	7.8%
Overweight (BMI 25-30)	150	30.6%
Obese (BMI >30)	302	61.6%
B/P (Systolic) mmHg <130	64.0	13.1%
>130	426.0	86.9%
Total Cholesterol (mmol/L) <4.0	95	19.2%
>4.0	395	80.8%
Triglyceride (mmol/L) <1.7	234	48.5%
>1.7	256	51.5%
Fasting Blood Sugar (mmol/L) <10	67	10.6%
>10	423	89.4%





Study Time: 1
MRN: K

TID	1.27	
LHR	0.26	
SSS 22	SRs 2	SDS 20
SS% 32	SR% 3	SD% 29

Proc ID	Persantin Stress:MBI
View ID	SAX-STR
Date	2013-12-08 13:39:10
Database	FemaleStressMB
Volume	85ml
Wall	152ml
Defect	57ml
Extent	38%
TPD	30%
Shape	0.61 [SI], 0.80 [Ecc]

Proc ID	Persantin Stress:MBI
View ID	SAX-RST
Date	2013-12-08 15:18:24
Database	FemaleRestMB
Volume	67ml
Wall	134ml
Defect	2ml
Extent	2%
TPD	1%
Shape	0.68 [SI], 0.76 [Ecc]

Str

Rst

Rest

Auto 0 - Grid Accept Reject



Figure 1: SPECT (Figure 1a): Stress induced ischemia (Arrows): anterior (white), apex (yellow) and antero-lateral wall (Green) shown on polar map (Figure 1b).

Semi quantitative data (Figure 1c): SSS = 22; SDS = 20 and Gated data (Figure 1d): LVEF (66% at rest, 57% at stress).

Coronary angiography (not shown): 75% stenosis of left anterior descending coronary artery and 60% left circumflex

The presence or absence of ischemia reported, as well as its location in the left ventricular (LV) wall. To assess the severity of ischemia several parameters were derived from C-equal images in (Figure 1) of a 58-year-old male patient with DM, hypertension and dyslipidemia referred for persantin MPI to rule out stress induced ischemia): the summed difference score (SDS) was used. The SDS is the difference between two semi-quantitative scores for 20 LV wall segments: the summed stress score (SSS), which assesses the perfusion defect during stress with respect to severity and extent, and the summed rest score (SRS), which assesses the extent and severity of the perfusion defect during rest. This measure used to describe the degree to which the deficit/ischemia is reversible. Based on SSS the MPI was considered normal with value of (0-3), mild abnormal (4–8), moderate (9–13), and severe (>13). [Reversible perfusion defect is defined as SDS \geq 2, mild stress induced ischemia (2–4), moderate (5–7), and severe (>7) [20].

In addition, the severity of ischemia assessed by the presence of transient ischemic dilatation (TID) with normal value of 1.22 and lung uptake expressed by lung to heart ratio (LHR) with normal value of 0.44. The left ventricular ejection fractions (LVEFs) at rest and stress also reported. All these parameters shown in Figure 1 as well.

Both qualitative and quantitative interpretation of imaging data carried by an expert in nuclear cardiology.

The final diagnosis of coronary artery disease based on coronary angiography according to a predefined protocol and interpreted by an experienced cardiologist.

The MPI findings shown in (Table 2). Our results showed the prevalence of silent myocardial ischemia in asymptomatic Saudi patients with type 2 diabetes mellitus was 21% and the positive predictive value of our MPI screening test 21%.

Table 2: Patient characteristics of the study group

Myocardial perfusion findings	Total (n=490)
Myocardial ischemia	237 (48.4%)
Coronary angiography	158 (66.7%)
Positive coronary angiography (coronary artery stenosis \geq 50%)	33 (21%)
SSS	5.60 \pm 10
SRS	4.1 \pm 8.9
SDS	1.7 \pm 2.9
TID	2 (1.0%)
LHR	0.3 \pm 0.1
LVEF (rest), mean \pm SD	56.2 \pm 13.3
LVEF (stress), mean \pm SD	56.5 \pm 14.1
LVEF difference, mean \pm SD	-0.4 \pm 15.9

Data are presented as n (%) and mean \pm SD

Abbreviations: SSS, summed stress score; SRS, summed rest score; SDS, summed difference score; TID, transient ischemic dilation; LHR; lung hear ratio, LVEF, left ventricular ejection fraction

Discussion

Patients with diabetes mellitus, in particular those with DM type 2 reported to be 2-4 times at higher risk of cardiovascular events when compared to non-diabetic partners [10,11, 21].

Stress radionuclide MPI used widely to evaluate patients with suspected or known ischemic heart disease. Several investigators have used MPI in the evaluation of patients with DM

and reported high prognostic values [22-25]. Bax. J et al reported that MPI was valuable test for indentifying asymptomatic diabetic patient who might require early and aggressive intervention to manage their cardiovascular risk [21]. However, one important question needed to be answered and that which diabetic patients needed to be screened for coronary artery disease and which test to use? To answer that question Wackers et al from Yale University conducted a large multicenter study for the detection of ischemia in asymptomatic diabetics (DIAD) trial included 1123

participants with well-controlled type 2 DM and they had no symptoms suggestive of coronary artery disease. They found that the rate of silent ischemia on stress MPI was 22% [14].

Detection of ischemia in asymptomatic diabetic Saudi patients (DIADS) was the aim of this study conducted on patients attending a large diabetic center. None of them had symptoms of coronary artery disease and assumed a good sample of asymptomatic Saudi patients with DM type 2. Our results showed that more than 20% of those older than 50 years have silent ischemia correlates very well with the original DIAD study by Wackers, *et al.* [14]. However, that was unpredictable considering the high prevalence of type 2 DM among Saudi population. The optimal medical therapy in the intervening period between the decision to refer to MPI and the test date could be the cause of low prevalence of abnormal MPI in our cohort study. If that assumption is accepted then a higher prevalence rate of silent ischemia in both controlled and non-controlled type 2 DM expected in a future large national study.

Conclusion

The prevalence of silent ischemia among asymptomatic Saudi patients with type 2 DM found to be 21%, which means that more than 2 out of 10 patients had silent myocardial ischemia. However and in order to justify the use of stress MPI as a screening test in this group of patients, a prospective large national study including both controlled and non-controlled type 2 DM is required.

References

1. Grundy SM, Benjamin IJ, Burke GL (1999) Diabetes and cardiovascular disease: a statement for healthcare professionals from the American Heart Association. *Circulation* 100: 1134-46.
2. Fuller JH (1993) Mortality trends and causes of death in diabetic patients. *Diabet Metab* 19: 96-9.
3. Al-Nozha MM, Al-Maatouq MA, Al-Mazrou YY (2004) Diabetes mellitus in Saudi Arabia. *Saudi medical J* 25: 1603-10.
4. Goraya TY, Leibson CL, Palumbo PJ (2002) Coronary atherosclerosis in diabetes mellitus: a population-based autopsy study. *Journal of the American College of Cardiology* 40: 946-53.
5. Donahoe SM, Stewart GC, McCabe CH (2007) Diabetes and mortality following acute coronary syndromes *JAMA* 298: 765-75.
6. Smith SC, Faxon D, Cascio W (2000) Diabetes and cardiovascular disease writing group VI: revascularization in diabetic patients. *Circulation* 105: e165-e169.
7. Deanfield JE, Maseri A, Selwyn AP, Ribeiro P, Chierchia S, et al. (1983) Myocardial ischemia during daily life in patients with stable angina: its relation to symptoms and heart rate changes. *Lancet* 2: 753-58.
8. Alexander XM, Landsman PB, Teutsch SM (2000) Diabetes mellitus, impaired fasting glucose, atherosclerotic risk factors, and prevalence of coronary heart disease. *Am J Cardiol* 86: 897-902.
9. Kang X, Berman DS, Lewin H (1999) Comparative ability of myocardial perfusion single-photon emission computed tomography to detect coronary artery disease in patients with and without diabetes mellitus. *American heart J* 137: 949-57.
10. Yamasaki Y, Nakajima K, Kusuoka H, Izumi T, Kashiwagi A, et al. (2010) Prognostic value of gated myocardial perfusion imaging for asymptomatic patients with type 2 diabetes: the J-ACCESS 2 investigation. *Diabetes Care* 33: 2320-6.
11. Bax JJ, Inzucchi SE, Bonow RO, Schuijff JD, Freeman MR & Global Dialogue Group for the Evaluation of Cardiovascular Risk in Patients with Diabetes. (2007) Cardiac imaging for risk stratification in diabetes. *Diabetes Care* 30: 1295-304.
12. Salehi Y, Fard-Esfahani A, Fallahi B, Aghahosseini F, Beiki D, et al. (2014). The myocardial perfusion scintigraphy in asymptomatic diabetic patients. *Iran J Nucl Med* 23: 27-35.
13. DeLuca AJ, Saulle LN, Aronow WS, Ravipati G, Weiss MB (2005) Prevalence of silent myocardial ischemia in persons with diabetes mellitus or impaired glucose tolerance and association of hemoglobin A1c with prevalence of silent myocardial ischemia. *The Am J Cardiol* 95: 1472-4.
14. Wackers FJ, Young LH, Inzucchi SE (2004) Detection of silent myocardial ischemia in asymptomatic diabetic subjects: the DIAD study. *Diabetes care* 27: 1954-61.
15. Mariano-Goulart D (2007) Myocardial perfusion imaging and cardiac events in asymptomatic patients with diabetes. *Heart Metabolism*, 35: 1-4.
16. Shmendi A, Pirie F, Naidoo DP, Tlou B, Pilloy W, et al. (2014) Myocardial perfusion imaging for evaluation of suspected ischemia and its relationship with glycemic control in South African subjects with diabetes mellitus.
17. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy 7: 545-52.
19. Berman DS, Kiaat H, Friendman JD (1993) Separate acquisition rest thallium-201/stress technetium-99m sestamibi dual isotope myocardial perfusion single photon emission computed tomography; a clinical validation study. *J am Coll Cardiol* 22: 1.
20. Sharir T, Germano G, Kang X, Lewin HC, Miranda R, et al. (2006) Prediction of myocardial infarction versus cardiac death by gated myocardial perfusion SPECT: risk stratification by the amount of stress-induced ischemia and the poststress ejection fraction. *J Nucl Med* 42: 831-7.
21. Bax J, Bonow R, Tschope D, In Zucehis, Barret E (2006) The potential of myocardial perfusion scintigraphy for risk stratification of asymptomatic patient with type 2 Diabetes *J Am Coll Cardiol* 48: 757-60.
22. Klocke FJ, Baird MG, Batemar TM, Berman DS, Carabello BA, et al. (2003) AcE/AHA/ASNC guidelines for the clinical use of cardiac Radionuclide imaging: a report of the American College of Cardiology, American Heart Association *Circulation* 108: 1404-918

23. Strauss HW, Douglas DM, Wittry MD, Cerquiera MD, Garcia EV, et al. (2008) Procedure guideline for myocardial perfusion imaging 33 J Nucl Med Techn 36: 155-61.
24. Kang X, Berman DS, Lewin HC Cohen I, Friedman JD, Germano G, et al. (1999) Incremented prognostic value of myocardial perfusion single photon emission computed tomography in patients with diabetes mellitus. Am Heart J 138: 1025-32
25. Vanzetto G, Halimi S, Hammoud T, Fagret D, Benhamou PY, et al. (1999) Prediction of cardiovascular events in Clinically selected high risk NIDDM patients. Diabetes Care; 22: 19-26

Submit your manuscript to a JScholar journal and benefit from:

- ¶ Convenient online submission
- ¶ Rigorous peer review
- ¶ Immediate publication on acceptance
- ¶ Open access: articles freely available online
- ¶ High visibility within the field
- ¶ Better discount for your subsequent articles

Submit your manuscript at
<http://www.jscholaronline.org/submit-manuscript.php>