

Appendico-Ileal Knotting a Rare Cause of Small Bowel Obstruction: Case Report and Literature Review from Leku General Hospital, Ethiopia

Asfaw Kibret^{1*}, Thomas Ameno² and Muluken Asres³

¹Department of Emergency Surgery, Leku general Hospital, Ethiopia

²Department of Anesthesia, Leku general Hospital, Ethiopia

³Department of general surgery, Leku general Hospital, Ethiopia

***Corresponding Author:** Asfaw Kibret, Department of Emergency Surgery, Leku general Hospital, Ethiopia, E-mail: aserkibret@gmail.com

Received Date: October 17, 2025 **Accepted Date:** October 28, 2025 **Published Date:** October 31, 2025

Citation: Asfaw Kibret, Thomas Ameno, Muluken Asres (2025) Appendico-Ileal Knotting a Rare Cause Of Small Bowel Obstruction: Case Report And Literature Review From Leku General Hospital, Ethiopia. J Surg Proce Case Rep 7: 1-7

Abstract

Small bowel obstruction is the common cause of acute abdomen for which emergency surgical intervention is mandatory. Bowel knottings like ileosigmoid knotting, ileoileal knotting, ileocecal knotting and appendicoileal knottings are rare occurrences as a cause of obstruction. Among these appendicoileal knotting is by far the rarest cause of obstruction. Since its 1st report in 1901, there are case reports of this scenario but all associated it with the presence of appendicitis. We report a 28 yrs old female patient who presented with sign symptoms of small bowel obstruction for whom appendicoileal knotting is identified with healthy looking appendix rapped around the distal ileum which was a near miss bowel segment which later returned healthy after the appendix release and warm saline soaked sterile pack rapped for a minutes. Appendicoileal knotting can occur in the absence of preceding appendicitis.

Keywords: Obstruction; Appendicitis; Appendicoileal Knotting; Double Loop Obstruction; Intraoperative Surprise



© 2025. Asfaw Kibret, Thomas Ameno, Muluken Asres. This is an open access article published by JScholar Publishers and distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Bowel obstruction in general is common cause for acute abdomen which most of the time needs emergency surgical intervention. Small bowel obstruction is the commonest cause worldwide. There are a lot of causes of small bowel obstruction [1, 2]. Among these, post-operative adhesion, hernia and malignancies are the most commonest causes to mention worldwide. But small bowel volvulus is the commonest in most part of our country Ethiopia. Bowel knotting like, ileosigmoid, ileoileal, ileocecal and appendicoileal are rare causes of bowel obstruction [12]. Among these appendicoileal knotting is the rarest to be occurred in bowel obstruction [3, 4]. Since 1901 of the 1st documented report there are few handful of cases reported in literatures [13]. Appendicoileal knotting is rapping of the appendix over the segment of distal ileum which causes closed loop obstruction. If untreated or delayed, it results in bowel vascular compromise and eventual bowel ischemia. Preoperative diagnosis is challenging since it has no pathognomonic features but mimic other intestinal obstruction causes of clinical presentations. Preoperative imaging, including computed tomography (CT), has shown potential utility, but its accuracy is limited and in resource limited setups its accessibility is ambitious [6, 7].

This case report adds another literature asset to the existing handful of cases world-wide with some peculiarities of uninflamed appendix causing appendico-ileal knotting in 28 yrs old female patient.

Case Presentation

This is a 28 yrs old female patient who presented to emergency room with the complain of abdominal pain of 3 days duration. She complained that she has crampy abdominal pain which started at the periumbilical area. In associated to this she had repeated episode of bilious type of vomiting, failure to pass feces and flatus. For this complain

she had been taken to her nearby private clinic and analgesics were given. But inspite of this her complain worsened and abdominal distension also started. She has no previous medical and surgical histories. She has 3 children and currently she is using family planning method. She had no vaginal bleeding, no vaginal discharge and no history of trauma.

Upon presentation she was acute sick looking with BP= 90/60 mmgh, PR=112 Bpm of tachycardia, RR=21 Breath/min, T=36.9, PSO₂= 96% off oxygen. On physical examination there was dry bucal mucosa, dry tongue and on abdominal examination there was abdominal distension with hypertympanicity on percussion, there was also minimal tenderness upon palpation. Digital rectal examination revealed empty rectum. Labratory investigation came with CBC= 16000, Hgb = 12g/dl, PLT=250000, RBS =150, Imaging suggested with multiple air fluid levels with preoperative diagnosis of SBO secondary to small bowel volvulus. For this diagnosis patient prepared for exploratory laparotomy. Medline abdominal incision used to open the abdominal cavity. Upon entry to the abdominal cavity there was moderately hemorrhagic free peritoneal fluid comes out and there was multiple small bowel loops distended, near to ischemia, especially the distal ileal segment. Surprisingly the was long non inflamed appendix rapped around the distal segment of the ileum near to the ileocecal junction, the tip of the appendix was buried to the ileal mesentery otherwise it was intact and no fecolith in it (Figure 1). Then the appendix released from around the entrapped distal ileal segment which was in double segment obstruction. Then after appendectomy done, the discoloured ileal segment rapped by warm saline soaked sterile surgical pack and time taken. Later the peristalsis and colour of the enterapped bowel segment returned normal. After meticulous observation of all segment of the small bowel, colon and other solid organ, no pathology witnessed then abdomen closed in layer and patient awaken and left OT stable. Her post operation course was uneventful, she discharged home on the 4th day and came 2 weeks later for follow in stable condition.

*Summarized tables of few case reports of appendicoileal knotting

Author And Year	Age	Sex	Clinical Presentatio	Intraoperative Findings	Procedure	Outcome
Mandal et al (2025) [1]	64	male	Abdominal pain, vomiting, inability to pass stool and flatus.	Appendix mucocele knotting causing cecal and ileal gangrene	Limited right hemicolectomy and double-barrel stoma.	Uneventful postoperative recover
Abule T,Chebo T,Billoro BB, (2022) [2]	30	Female	Colicky abdominal pain, vomiting, and constipation	Appendico-ileal knot causing small bowel obstruction (SBO).	Untwisting of the knot, appendectomy	Uneventful recovery, discharged on day 4.
Idowu NA et al. (2024) [3]	72	Male	Colicky abdominal pain, nausea, vomiting, constipation, and fever over 4 days.	Appendico-ileal knotting with gangrenous appendix and terminal ileum.	Limited right hemicolectomy and ileo-colonic anastomosis.	Discharged on postoperative day 10.
Kifle T et al (2018) [4]	46	female	Acute abdomen with severe pain, bilious vomiting, and constipation for 7 days.	Confirmed appendiceal knotting with apperndicular mucocele causing SBO.	Laparotomy with untwisting of the knot, followed by appendectomy.	Recovered well.and discharged on the 5 th post op date
Kabuye U et al. (2024) [5]	28	female	Abdominal pain, vomiting, and constipation. Initial diagnosis was SBO from an intestinal band.	Appendix entangled around the terminal ileum, causing a closed-loop obstruction. The appendix was gangrenous, but the ileum was viable.	Retrograde appendectomy after releasing the knot.	Uneventful recovery, discharged on day 4.
Zewdu D et al. (2022)[6]	34	male	SBO symptoms	Ileoappendicular knotting	Appendectomy,with resection of gangrenous bowel	Full recovery, though experienced post-op diarrhea.
Kassahun B et al. (2025)[7]	28	male	Progressive lower abdominal pain, nausea, anorexia, and bilious vomiting for 2 days.	Gangrenous 20 cm appendix tightly encircling 1 m of gangrenous distal ileum.	Untwisting of the knot, appendectomy, resection of the gangrenous ileum, and end-to-side ileo-transverse anastomosis.	Uneventful recovery, discharged on postoperative day 6.
Klein et al. (2024)[8]	80	Female	Small bowel obstruction sign and symptom of 1 day duration	Appendicular knot of healthy appendix,obstructing the small bowel as closed loop obstruction	appendectomy	Discharged well
Alemu et al. (2023)[9]	50	Female	Abdominal pain, vomiting, abdominal distension.	Appendicoileal knotting with gangrenous ileum and appendix mucocele	Bowel and appendix resection with ileotransverse anastomosis	Discharged on the 7 th post-operative date
Lin, Tso-Lin et al. (2017)[10]	4	male	Vomiting,abdominal pain	Inflamed appendix causing appendiculoileal band knot	appendectomy	Discharged uneventfully



Figure 1: Intraoperative Finding of Appendicoileal Knotting Revealing Appendix Rapped Around the Distal Ileal Segment

Discussion

Appendico-ileal knotting remains an exceedingly rare cause of small bowel obstruction (SBO), often presenting a significant diagnostic challenge. As evidenced by the collected case reports, the condition lacks a pathognomonic clinical presentation, with symptoms typically overlapping with other, more common, causes of SBO, such as volvulus, adhesions or hernias. This leads to most diagnoses being made intra-operatively, often as a "surprise" finding during exploratory laparotomy [15, 16].

The patient demographic in these cases varies widely, spanning from pediatric patients to the elderly. The

specific pathophysiology depends on an elongated, mobile appendix, which can become inflamed and form a constricting band around a loop of the ileum. The outcome is critically dependent on the timing of surgical intervention. Early intervention, as seen in cases where the bowel remains viable, allows for a straightforward procedure involving the untwisting of the knot and appendectomy. In contrast, delayed presentation or diagnosis, especially in resource-limited settings, can lead to catastrophic complications such as bowel gangrene, perforation, and septic shock, requiring more extensive and complex procedures, such as bowel resection and anastomosis [1-10].

Histopathological findings, when available, some-

times reveal an associated mucocele of the appendix [1, 9]. These findings suggested that certain appendiceal pathologies might contribute to the knotting mechanism. In most cases appendicitis was mentioned as a preceding incident to appendicoileal knotting. Yet, there is one report that reveal the occurrence of appendicoileal knotting in the presence of macroscopically and microscopically healthy looking appendix [8]. Our case also presented with the same finding with this, supporting the occurrence of appendicoileal knotting in the absence of preceding appendicitis. So appendix can be a cause for small bowel obstruction in the presence of its inflammation or only mechanically as a band in the absence of its inflammation. There are two basic situations where the appendix may also cause a mechanical obstruction appendicular tip attached to the mesentery surrounding an ileal loop, producing compression of its lumen and the appendicular tip attached to the intestinal serosa, producing the obstruction by direct compression or torsion of a loop. The overall literature emphasizes that while appendico-ileal knotting is a rare event, a high index of clinical suspicion is necessary in patients presenting with SBO, especially in the absence of a clear etiology like a history of prior surgery. Early diagnosis and prompt surgical management are the cornerstones of successful treatment and significantly improve patient outcomes. But the overall management depends on the viability of the bowel or strangulation [11, 14, 15].

Conclusion

Appendico ileal knotting is still rare cause for small bowel obstruction. But as abdomen is “a Pandora box” the very rare things can happen and clinical suspicion

is needed. Early diagnosis may help the patient for early surgical intervention. Appendico ileal knotting can occur in the absence of appendicitis.

Acknowledgement

We would like to thank all leku general hospital clinical staffs who are involved in the management of the patient specially to the operating theater nurses and anesthetists.

Fund

No fund for this case report.

Informed Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Ethical approval

N/A

Conflict of Interest

No conflict of interest.

Data Availability

All relevant data are within the paper and its Supporting Information files.

References

1. Mandal M K, Sarkar N, Sarkar S, & Mollah J (2025) A case of appendiculo-ileal knotting as a cause of gangrenous small bowel obstruction in a 64-year-old man in a Government Hospital, West Bengal, India. *International Surgery Journal*. 12: 1383–86.
2. Abule T, Chebo T, Billoro BB (2022) Appendico-ileal knotting causing small bowel obstruction: A case report. *Clin Case Rep*. 10: e05878.
3. Najeem Adedamola IDOWU, Waheed Olalekan IS-MAEEL, et al. (2024) Appendico-Ileal Knotting: A Rare Cause of Strangulated Small Bowel Obstruction. *Ethiopian Journal of Health Sciences*. 34.
4. Kifle Anteneh Tadesse, Tesfaye Samuel (2018) Appendico ilial knotting: a rare cause of small bowel obstruction. *J. Surg. Case Rep*. 2018.
5. U Kabuye, J Damulira, MD Okuku (2024) Appendico-ileal knot: a rare form of small bowel obstruction: a case report *Int.J.Surg. Case Rep*: 123.
6. Wondwosen M, Tantu T, Zewdu D (2022) Appendiceal knotting causing small bowel obstruction: A rare case report. *International Journal of Surgery Case Reports*. 93: 106970.
7. Berhanu Kassahun, Abebe Dilie Afenigus (2025) Gangrenous small bowel obstruction resulting from appendico-ileal knotting: a rare case report from Bete Markos Medical and Surgical Center, Ethiopia, *Journal of Surgical Case Reports*. 539.
8. Klein Nulend R, Quinn R, Fok KY, Pathmanathan N (2024) The appendix: An unexpected band obstruction. *J Case Rep Article ID*: 100129Z12RN2024
9. Alemu DS, Arefayne M, Bekele T, et al. (2023) A Case of Appendiculo-Ileal Knotting As a Cause of Gangrenous Small Bowel Obstruction in a 50-Yr-Old-Woman, in a Private Hospital. *J Surg Surgic Case Rep*. 4: 1036
10. Intestinal obstruction caused by appendiceal knot Lin, Tso-Lin, et al. (2019) *Pediatrics & Neonatology*, Volume 60: 216–17.
11. Bhandari L, Mohandas P (2009) Appendicitis as a cause of intestinal strangulation: a case report and review. *World J. Emerg Surg*. 4: 34.
12. Kibret A (2023) Ileo-Ileal Knotting a Rare Cause of Double Loop Small Bowel Obstruction, Diagnostic and Intra-operative Challenge at Ethiopian Leku District Hospital, Sidaama Region : Case Report. *Clin Surg*. 8: 3657.
13. Hotchkiss LW (1901) Acute intestinal obstruction following appendicitis: A report of three cases successfully operated upon. *Ann Surg*. 34: 660–77.
14. Nulend RK, Quinn R, Fok KY, Pathmanathan N (2024) The Appendix: An Unexpected Band Obstruction. *Journal of Case Reports and Images in Surgery*. 10: 1-9.
15. JG M, SA K, LJ A, EA A (2017) Intestinal obstruction caused by appendicitis: a systematic review. *Journal of the West African College of Surgeons*. 7: 94.
16. Awale L, Joshi BR, Rajbanshi S, Adhikary S (2015) Appendiceal tie syndrome: A very rare complication of a common disease. *World J SGastrointest Surg*. 7: 67–70.

Submit your manuscript to a JScholar journal and benefit from:

- § Convenient online submission
- § Rigorous peer review
- § Immediate publication on acceptance
- § Open access: articles freely available online
- § High visibility within the field
- § Retaining the copyright to your article

Submit your manuscript at
<http://www.jscholaronline.org/submit-manuscript.php>